

BIOMIN[®] N



5-0-0

Product Information

BIOMIN N is a liquid plant nutrient product derived from hydrolyzed soy protein. It is designed for soil and/or foliar applications.

GUARANTEED ANALYSIS

Total Nitrogen (N).....5.0%
5% Water Soluble Nitrogen

Derived from soy protein hydrolysate.

F1416

KEEP OUT OF REACH OF CHILDREN

**SHAKE WELL BEFORE USE
DO NOT APPLY AT TIME OF BLOOM**

NET CONTENTS: 2.5 gal. (9.45 l)
Weight Per Gallon: 8.5 lb. (3.8 kg) @ 68°F
Lot #:



Made in USA



Directions and General Recommendations

Dilute **BIOMIN N** with at least 200 parts water and apply in accordance with the directions.

VEGETABLES, POTATOES, SUGARBEEF, ROW CROPS, AND ROOT CROPS: 1-3 gal./ac. at 2 week intervals after thinning or when plants are well established.

COMMERCIAL GREENHOUSES AND NURSERIES: Inject into irrigation water after filter, at ratio of at least 1:1,000. Apply at 2 week intervals.

ALFALFA: 1-3 gal./ac. one week after each cutting.

ALMONDS, APRICOTS, CHERRIES, HOPS, PLUMBS, PEACHES, GRAPES, APPLES, PECANS, PISTACHIOS, PRUNES, WALNUTS, AND OTHER FRUITS AND NUTS: 2-3 gal./ac. prior to bloom. Older trees may require another application 4-6 weeks later. Repeat the application 2 more times in 3-4 week intervals.

BEANS, PEAS, SOYBEANS, TOMATOES, PEPPERS: 1-3 gal./ac. at bud stage, prior to bloom. Repeat the application 2 more times in 3-4 week intervals.

CITRUS, AVOCADOS: 3-5 gal./ac. at bud stage, prior to bloom. When new growth appears, repeat the application 2 times at 2-4 week intervals.

COTTON: 2-3 gal./ac. after true leaves appear. Second application should be made at early square at 3-4 gal./ac. Repeat the application 2 more times at 3-4 week intervals.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

RICE, BARLEY, SORGHUM, WHEAT, AND OTHER SMALL GRAINS: 2-3 gal./ac. at boot stage. Repeat the application prior to and after bloom.

STRAWBERRY AND OTHER BERRIES: 1-3 gal./ac. after transplanting. Repeat the application throughout the season at 2-4 week intervals.

POTATOES: 1-3 gals/ac. at tuber initiation. Repeat the application at 2-4 week intervals.

CORN: 1-3 gal./ac. prior to tasseling. Repeat the application at 2-4 week intervals.

SUNFLOWER AND OTHER OILSEED CROPS: 1-3 gal./ac. when plants are well established. Repeat the application at 2-4 week intervals.

BIOMIN N 5-0-0 is recommended for use on all crops including: alfalfa, almond, apple, apricot, avocado, barley, beans, carrot, cherry, citrus, corn, cotton, grape, hops, peach, pear, pecan, pepper, pistachio, plum, potato, prune, rice, row crops, sorghum, soybean, strawberry, sugarbeet, sunflower, tomato, vegetable, walnut, and wheat.

Precautionary statement: do not swallow. Avoid contact with eyes. **First Aid:** if swallowed: seek medical attention. Rinse mouth. If in eyes: remove contact lenses if present. Wash thoroughly with water for several minutes. **Storage & Disposal:** store in a safe manner. Dispose product/container in accordance with federal, state, and local regulations.

JHBiotech, Inc.

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006 U.S.A. 805-650-8933 jhbiotech.com

Copyright © 2016 All rights reserved

0342-D-B-1